

Mapping inequities in global vaccine sentiment research

Duilio Balsamo,^{1,2} Vittoria Offeddu ,¹ Zhina Aghamohammadi,¹ Chiara Chiavenna,^{1,2} Laura Pasqua Leone,^{1,3} Elena D'Agnese,^{1,4,5} Deepak Sharma,⁶ Aleksandra Torbica,^{1,7} Soheil Shayegh ,^{8,9} Javier Andreu-Perez,^{10,11} Alessia Melegaro^{1,2}

To cite: Balsamo D, Offeddu V, Aghamohammadi Z, *et al*. Mapping inequities in global vaccine sentiment research. *BMJ Glob Health* 2025;**10**:e018587. doi:10.1136/bmjgh-2024-018587

Handling editor Helen J Surana

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjgh-2024-018587>).

DB and VO are joint first authors.

Received 10 December 2024

Accepted 25 September 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Alessia Melegaro;
alessia.melegaro@unibocconi.it

ABSTRACT

Introduction Negative public sentiment towards vaccination (PSV) poses significant challenges to the effectiveness of immunisation programmes, with dramatic effects on morbidity and mortality for vaccine-preventable diseases. Yet, health research is often shaped by economic and geopolitical factors rather than countries' epidemiological or healthcare needs. This study examines global patterns and drivers of PSV research on five vaccines—polio, measles, human papillomavirus, influenza and SARS-CoV-2—and evaluates the COVID-19 pandemic's impact on research volume, focus and distribution.

Methods We conducted a machine-learning-assisted literature search on PSV without geographical, language or time constraints. Using natural language processing, network and statistical analyses, we examined the global PSV research landscape and identified geographical, epidemiological and economic drivers.

Results We analysed 13 287 articles and detected consistent literature growth from 1980 onwards, with vaccine-specific peaks following key licensing events. After 2020, publication volumes rose above projections for influenza (32%; 95% CI 20% to 46%) but declined for polio (−56%; 95% CI −68% to −26%) and measles (−17%; 95% CI −33% to 9%). Although PSV research had global coverage, its distribution was markedly imbalanced, largely shaped by country-specific economic factors. A few high-income countries (HICs) produced 72% of publications and the likelihood of a country being studied varied by income and vaccine. Foreign authorship also increased as the income of the studied country decreased (over 75% in low-income and middle-income countries vs below 50% in HIC).

Conclusions PSV research reveals persistent inequities, with a misalignment between countries leading research and populations most in need of its outcomes. These inequities, further exacerbated by COVID-19 disruptions, reflect systematic imbalances in global health. Our findings underscore the need to decolonise research by fostering leadership, agenda-setting and accountability that centre on the necessities of affected communities. Achieving this will require funding and publication reforms that promote equitable collaborations and elevate local priorities alongside long-standing global health objectives.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Public sentiment towards vaccination (PSV) significantly influences vaccination campaign outcomes. While bibliometric studies have explored PSV research patterns, economic and contextual drivers in this research field remain underexplored, and international collaboration dynamics lack in-depth analysis.

WHAT THIS STUDY ADDS

⇒ This study identifies significant economic disparities shaping PSV research, with high-income countries dominating the field and directing studies towards low-income settings with limited local author representation. It also highlights the COVID-19 pandemic's role in shifting the research focus from critical childhood vaccines to SARS-CoV-2, exacerbating these imbalances.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ By revealing inequities in PSV research, this work underscores the need for more equitable and inclusive research initiatives, prioritising public health needs while addressing systemic disparities in global health research and policy.

INTRODUCTION

The primary goal of global health research is to promote health equity and strengthen healthcare systems by bridging global knowledge with local interventions.¹ However, global health initiatives are heavily influenced by economic factors.² Researchers from low- and middle-income countries (LMICs) face limited access to global health research grants³ and are under-represented in peer-reviewed publications.⁴ Furthermore, 90% of all health research funding targets health issues of industrialised countries,⁵ marginalising lower-income regions, where health burdens are typically greater. This perpetuates inequities, with researchers from high-income countries (HICs) dominating research activities also in

LMICs, resulting in inadequate research agendas that fail to meet local public health needs.² Vaccine uptake and childhood immunisation patterns across the globe represent a significant example of this dynamic.

Despite vaccines' efficacy, global uptake remains suboptimal in most countries, pockets of zero-dose individuals persist,⁶ and public sentiment towards vaccination (PSV) is imbued with widespread hesitancy.⁷ Over time, considerable research efforts have been directed towards investigating the intricate mechanisms shaping PSV, resulting in a heterogeneous, multidisciplinary and fast-growing body of literature.⁸ Numerous reviews attempted to systematise and summarise the existing evidence on PSV.⁹ However, due to the sheer amount and complexity of the published data, this task is proving increasingly difficult, especially without the aid of machine-assisted methods. Identifying the main drivers of PSV research might facilitate the definition of future research priorities and provide an evidence-based rationale to support a more appropriate allocation of resources.

To what extent PSV research is shaped by the interplay between economic factors, local disease endemicity and the effective need to study PSV locally is not fully understood,^{8 10–12} especially in light of disruptive events, such as the COVID-19 pandemic. The exceptional circumstances in which the SARS-CoV-2 vaccine was licensed, together with the determining role it has played in curbing the pandemic, significantly intensified research efforts and public discussions on PSV.¹³ However, the pandemic also diverted a significant portion of global academic efforts and funding from other research to COVID-19-related contributions, which benefitted from fast publication tracks and great academic attention, putting research on other diseases and vaccines at considerable disadvantage.^{14–16}

In this work, we investigated the global patterns and drivers of scientific research on PSV, including studies on public attitudes, beliefs, emotions, trust, hesitancy and uptake. To enable a fine-grained yet broad perspective on the literature, we used machine-learning-assisted article selection to perform a large-scale literature search of peer-reviewed publications. We screened for a selected set of vaccines, *i.e.* those against polio, measles, human papillomavirus (HPV), influenza and SARS-CoV-2, chosen to be representative of different target populations, licensing and distribution histories, and viral pathogens with distinct transmission routes and epidemiology. Using a combination of natural language processing and large language model techniques, we enriched each record with standardised geographic metadata about authorship and place of study and linked each record with epidemiological and economic country-specific metadata. We analysed publication trends for all vaccines, and by benchmarking publication volumes before and after 2020, we quantified the extent to which the COVID-19 pandemic has altered pre-existing publication dynamics. Then, leveraging network and statistical tools, we modelled and analysed the geographical, epidemiological and

economic factors driving scientific focus and academic contribution in the PSV research field. The integrated perspective introduced in this work charts the overall structure of PSV literature at large scale, allowing for direct comparisons across a set of vaccines with distinct characteristics and public attention patterns.

Our paper represents a significant contribution to the literature both in terms of methodology as well as implications for academic research and global health policy. By combining novel techniques with traditional statistical approaches, we offer a strong example of how diverse methods can be jointly applied to address complex research questions. By outlining the structural inequities that influence PSV research agendas, our findings provide important insights to guide future research priorities, with the ultimate goal of a more equitable, balanced and resilient global health research agenda.

METHODS

Database construction

To retrieve scientific publications on PSV, we developed a comprehensive search string according to validated guidelines,¹⁷ incorporating keywords regarding public knowledge, attitudes, awareness, emotions and beliefs related to vaccination, vaccine hesitancy and refusal, trust and uptake (online supplemental file 1). To optimise the specificity of our search, we limited vaccine-related keywords to the article title, and other search terms to the title and abstract. We refined our search string through subsequent search cycles and qualitative screening of the first $n=100$ delivered papers, iteratively removing (1) keywords retrieving irrelevant literature, *e.g.* 'vaccinia', and (2) individual terms that were likely to occur together with a synonymous search term in true positive cases and retrieving irrelevant literature otherwise. The literature search was conducted in the MEDLINE and Web of Science Core Collection databases until December 2023, without geographical, language or other time constraints. We chose this database combination to cover a broad spectrum of scientific disciplines while obtaining the best possible balance between sensitivity and specificity.¹⁸ We removed duplicates using the article DOIs and titles and excluded papers not mentioning any of the vaccines of interest in their titles or abstracts.

Given the large number of retrieved articles, we employed supervised machine learning (ML) techniques to identify studies directly related to PSV. We began by manually annotating a sample of 1000 articles by title and abstract, following a predefined set of inclusion criteria. Articles were retained if they addressed at least one dimension of vaccine sentiment, such as studies on knowledge, attitudes and practices, or those that assessed determinants of vaccine acceptance, uptake or timeliness. Eligible study types included both quantitative and qualitative primary research, secondary data analyses, theoretical frameworks and mathematical models exploring vaccination determinants, as well as reviews and

editorials. After the first round of screening, performed by two independent reviewers, we obtained an inter-rater agreement Cohen's $k=0.8$, indicating substantial agreement. A third reviewer resolved the disagreements, resulting in 376 articles labelled as directly related to PSV and 624 as unrelated.

Next, using this annotated sample, we trained multiple ML classifiers, fine-tuning the pipelines' hyperparameters in a fivefold cross-validation setting to maximise classification performance. Model evaluation was conducted on a held-out test set (30% of the annotated data), achieving high performance across models, with all classifiers reaching an F1-score >0.86 and a Matthews correlation coefficient (MCC) >0.780 .¹⁹ The best-performing model—Random Forest Classifier with bag-of-words and tf-idf preprocessing—achieved an F1-score of 0.89, and MCC of 0.82 on the test set. This model was then applied to classify the full dataset. For each retained article, we identified the primary vaccine studied, defined as the most frequently mentioned vaccine in the title and/or abstract, as well as the other mentioned vaccines. If multiple vaccines were referenced equally, the first vaccine appearing in the text was assigned as the primary one.

Data preprocessing and enrichment

Using OpenAI's GPT (online supplemental file 1), we enriched our corpus by extracting from titles and abstracts of collected articles the authors' countries, *i.e.* the countries of the authors' institutions, and the studied countries, *i.e.* the countries of the analysis. Each country was then assigned a list of categorical attributes, including income level as of 2021,²⁰ time-dependent vaccine-specific coverage^{21 22} and time-dependent disease incidence for each related disease.^{23 24} Country income was chosen as a proxy for a country's research capacity since research institutions, such as universities and public health organisations, largely rely on national funding and infrastructure to conduct their projects. Disease incidence dynamically captures the epidemiological pressure that a country is facing. Together with varying levels of vaccination coverage, these factors are likely to influence the scientific and public concern about a health threat, potentially fuelling or slowing down research. All attributes were categorised into four groups (L=low; LM=lower-middle; UM=upper-middle; H=high) based on the empirical quantiles (online supplemental file 1). Polio incidence was classified as either 'no cases' or 'reported cases'. Due to incomplete data on disease incidence and vaccination coverage for HPV and influenza, we limited the analysis including all attributes to measles and polio.

Time trend analysis

To characterise the publication trends over time, we modelled the volume of published papers for each vaccine from 2010 to 2019 fitting an exponential function. We conducted this analysis starting from 2010 to

ensure inclusion of the period following the licensure of both the HPV and 2009 H1N1 pandemic influenza vaccines. The impact of the SARS-CoV-2 pandemic on PSV-related research was assessed by comparing the predicted publication volumes for 2020–2023, based on the fitted exponential models, with the actual volume produced after 2020, calculating the percentage difference (online supplemental file 1).

Geographical network construction

We developed a custom network analysis pipeline to map the geographic direction of global research on PSV. For each 5-year time interval, we created vaccine-specific networks where each node represents a country. Each directed weighted edge, a connection between countries i and j , was formed proportionally to the number of papers authored or coauthored by institutions from country i focusing on country j . Articles with multiple authors' countries and/or studied countries contributed to multiple edges in the network. We further characterised each node i by counting the number of (1) Domestic studies, papers on country i authored by local researchers alone (the weighted self-loops of node i); (2) Inbound studies, papers on country i authored or coauthored by foreign institutions (the weighted in-degree minus the self-loops) and (3) Outbound studies, papers on any foreign country authored or coauthored by institutions from country i (the out-degree minus the self-loops).

Statistical analysis on research drivers

We employed a regression framework to investigate the drivers of global PSV research. First, to investigate the probability of a country being studied, we fitted logistic regression models using a binary dependent variable indicating whether a country had been the object of at least one research paper on a given vaccine (1 if studied, 0 if not) (online supplemental file 1). Second, for countries that were studied, we analysed which country-level factors influenced the proportion of papers authored or coauthored by foreign institutions (inbound/(inbound+domestic)). We achieved that through a Beta regression model, suitable for modelling proportions (online supplemental file 1). Independent variables in both models included (1) time interval, classified as prepandemic (2010–2014; 2015–2019) and postpandemic (2020–2023) periods, (2) country income level, (3) country time-varying vaccination coverage level and (4) country time-varying disease incidence level. All models except the ones for polio (for which not enough data was available) included an interaction term between income and time to test whether the probability of being studied or the proportion of foreign authorship varied over time across income groups (online supplemental file 1).




Figure 1 Annual number of published scientific articles on public sentiment towards five selected vaccines (reported in different colours) from 1980 onwards. Dashed lines indicate the total number of articles mentioning each vaccine in their title and/or abstract. Solid lines indicate the total number of articles focusing on a specific vaccine as the primary vaccine. Vertical lines mark vaccine licencing dates. The licencing date for influenza corresponds to the vaccine against the 2009/H1N1 pandemic strain, while the licencing of the 1945 inactivated seasonal influenza vaccine is omitted. Polio (1955) and measles (1963) vaccine licencing dates also precede the emergence of PSV research. For visualisation purposes, the number of papers published on SARS-CoV-2 is evaluated and reported quarterly. HPV, human papillomavirus.

RESULTS

The unfolding of PSV research over time

Our query across two databases yielded 154552 articles from 1902 to 2023 (online supplemental figure S1). We removed duplicates (43%) and screened for references to any of the five selected vaccines, retaining 17587 articles. Of these, 97.5% were written in English, 2.2% in other European languages and 0.2% in non-European ones. We then used ML-assisted techniques to exclude non-relevant papers (24% of articles on the selected vaccines). We retained a total of 13287 articles (18% of the initial yield) for analysis (online supplemental figure S1). Of these, 44% focused on SARS-CoV-2 as the primary vaccine (n=5818), 25% on influenza (n=3328), 23% on HPV (n=3078), 7% on measles (n=883) and 1% on polio (n=180).

The volume of PSV literature did not rise significantly until the late 1980s (figure 1). We observed exponential growth across all vaccines until 2020, with vaccine-specific surges corresponding to the licencing of the HPV and SARS-CoV-2 vaccines, and a volume spike in influenza-related publications following the development of the novel influenza vaccine against the 2009/H1N1 pandemic strain. Using the fitted exponential models (online supplemental figure S2), we predicted the volume of publications post-SARS-CoV-2 and compared them against the measured volumes, finding a higher-than-expected increase in the volume of influenza-related research (perc. diff.=32%; 95% CI 20% to 46%). In contrast, the volume of PSV research on the two routine childhood vaccinations, polio and measles, fell below expectation after the pandemic, with decreases of perc. diff.=−56% (95% CI −68% to −26%) and perc. diff.=−17% (95% CI −33% to 9%), respectively.

Notably, an increasing proportion of papers mentioned more than one vaccine in the text (7.3% before 2020 and 12.9% after 2020). In particular (figure 2), vaccines sharing similar target populations, such as polio and measles, or those protecting against pathogens with similar characteristics, such as influenza and SARS-CoV-2, two respiratory viruses with pandemic potential, were more frequently studied together. Additionally, several studies on all primary vaccines began co-mentioning SARS-CoV-2 during the pandemic. This applied even to vaccines with relatively different characteristics, such as polio, measles or HPV, suggesting a potential impact of the SARS-CoV-2 pandemic on PSV research. Despite historical trends, the share of articles studying influenza alone also declined after the pandemic, with a large proportion (19.7%) of recent influenza-related articles also referencing the new coronavirus.

Geographical patterns in global health research

We mapped the geographical distribution of global PSV research from 2010 onwards (figure 3). To reconstruct the direction of research, we evaluated country-specific proportions of publications authored by local institutions alone (domestic), authored by or with institutions from a foreign country (inbound) or focused exclusively on foreign countries (outbound).

Overall, published PSV literature covered most countries worldwide, with HIC in North America and Europe authoring the majority of studies (72.2%). However, notable differences emerged in the spatial distribution of published articles across the examined vaccines. PSV research on the polio vaccine was more geographically clustered compared with other vaccines, with hotspots of research in sub-Saharan Africa, the Middle East and




Figure 2 Co-occurrence of primary vaccines and mentioned vaccines from 2010 onwards. For each 5-year interval, the colour-coded heatmap shows the percentage of articles focusing on a primary vaccine (y-axis) and mentioning another vaccine (x-axis). Percentages on the diagonal represent the proportion of papers that focus exclusively on a primary vaccine. Articles mentioning more than one additional vaccine are included more than once in the matrices. The total number of articles focusing on each primary vaccine for a given period is reported under the y-axis labels. HPV, human papillomavirus.

South Asia. Conversely, research efforts on the influenza vaccine were relatively frequent in Europe and North America but scarce on the African continent, with respectively >70%, >40% and <30% of countries involved, *i.e.* authoring or being studied in at least one paper ((online supplemental figure S3), top row).

We also observed income-related global imbalances in terms of research direction (online supplemental figure S3, bottom row). Despite the high number of studies on polio and measles vaccines authored by institutions in North America, Europe and Australia, the direction of these studies was predominantly outbound for HIC (median proportion outbound studies, high-income; polio: 82%; measles: 61%), and mostly inbound for low-income countries (median proportion inbound studies, low-income: polio: 100%; measles: 78%). In the same time frame, little domestic and even less outbound research emerged from low-income and lower-middle-income countries. Similar but reversed income-dependent trends were observed in the distribution of PSV research on the HPV and influenza vaccines, although in HIC the shares of domestic, inbound and outbound research efforts were more equally distributed for these two vaccines (median proportion domestic, inbound, outbound studies, high-income; influenza: 24%, 37%, 26%; HPV: 33%, 32%, 33%). Lastly, we measured comparable shares of outbound research on SARS-CoV-2 across income levels (median proportion outbound studies SARS-CoV-2: low-income: 21%; low-middle-income: 31%; upper-middle-income: 33%; high-income: 40%), with the amount of inbound research decreasing with increasing country income, mirroring patterns observed for other vaccines.

Drivers of global research flow

Through vaccine-specific logistic regression models, we assessed the potential factors influencing the probability of countries being the focus of PSV research. The reconstructed marginal probabilities are shown in figure 4 and online supplemental figure 4, upper panels.

Coefficients are reported in online supplemental tables S2 and S3. We found that a significantly lower proportion of countries has been studied in polio research compared with all the other vaccines. For all vaccines, the probability of being studied was significantly associated with the country's income level, although this dependence followed opposite directions for different vaccines. In the reference period (2010–2014), lower-income countries had a significantly higher probability of being the object of polio or measles PSV research compared with HIC. The disparity was greatest for the measles vaccine in the reference period, with HIC having 98% lower odds of being studied compared with the low-income group (H vs L: OR=0.02; 95% CI 0.00 to 0.11) (online supplemental table S2). This income disparity remained constant over time for polio and appeared to level up for measles, with increased interest in higher-income settings but a decrease in lower-income ones. In contrast, HIC had a significantly higher probability of serving as study locations for the HPV and influenza vaccines, perpetuating existing income disparities without significant time-and-income variation. Additionally, the probability of being studied for the influenza vaccine appeared to be significantly higher in the years following the COVID-19 pandemic (2020–23 vs 2010–14: OR=7.57; 95% CI 1.16 to 149.71) (online supplemental table S3).

For both measles and polio, local disease incidence affected the probability of countries to be studied (online supplemental figure S4, table S2). Countries where clinical polio cases were reported had 2.7 times higher odds of being studied compared with countries where the disease was eliminated (reported cases vs no cases: OR=2.67; 95% CI 0.95 to 7.37). Analogously, the probability of being studied was significantly higher among countries experiencing comparatively high measles incidence (H vs L: OR=10.11; 95% CI 4.45 to 24.76). Conversely, we did not detect a consistent effect of national vaccination




Figure 3 Geographical distribution of author countries and studied countries by vaccine from 2010 onwards. (a-e) chart different vaccines — Polio, Measles, HPV, Influenza, SARS-CoV-2— respectively, with (f–j) focussing on European countries. Pie slices represent country-specific shares of domestic, inbound or outbound studies. Domestic: studies on country *i* authored by local researchers alone. Inbound: studies on country *i* authored or coauthored by foreign institutions. Outbound: studies on any foreign country authored or coauthored by institutions from country *i*. The size of each pie chart is proportional to the total number of studies involving a country. Background colour corresponds to countries' income levels. H, high income; L, low income; LM, lower-middle income; UM, upper-middle income; HPV, human papillomavirus.

coverage on the probability of being studied for either vaccine.

Determinants of foreign authorship

For countries that have been the object of PSV research, we modelled the proportion of articles (co)authored by foreign countries using Beta regression models. The fitted values are shown in figure 4 and online supplemental figure S4, bottom panels. The coefficients are reported in online supplemental tables S2 and S3. Across all vaccines, during the reference period (2010–2014), the proportion of papers involving foreign research efforts increased as the income level of the studied country decreased. This income-related disparity appeared to diminish over time for the measles, HPV and influenza vaccines. Conversely, we observed no significant time-dependent change for polio research.

For both polio and measles, countries reporting clinical disease cases had a smaller proportion of articles (co) authored by foreign researchers. We found no consistent effect of national vaccination coverage on the proportion of foreign authorship.

DISCUSSION

This work examines the drivers of global health research using innovative methodologies, offering novel insights and analyses that, to our knowledge, are unexplored in the existing literature. We integrated an extensive literature search, ML-assisted article screening and data enrichment, network analysis and statistical methods to develop a novel database of research papers on PSV, which offers a valuable resource for addressing broad global health research questions. Going beyond traditional




Figure 4 (Top row) Estimated probabilities, with 95% CIs, of being the subject of PSV research on a given vaccine across time points and income groups. (Bottom row) Estimated proportion, with 95% CIs, of PSV research articles on a given vaccine involving foreign researchers (inbound/(inbound+domestic)), by income level of studied countries. H, high income; L, low income; LM, lower-middle income; HPV, human papillomavirus; PSV, public sentiment towards vaccination; UM, upper-middle income.

bibliometric methods^{8 10–12} and merging information on country-specific epidemiological and economic factors as well as direction of research, we investigated patterns and drivers shaping the global research efforts on PSV.

Our findings portrayed PSV research as a rather recent research field, partially shaped by vaccine-specific epidemiological conditions and unforeseen global health challenges, such as pandemic threats, but strongly influenced by country-specific economic determinants that steer academic focus and the opportunity for scientific collaborations.

By modelling the volumes of historical research data, we revealed that overall PSV publications have grown exponentially until 2020. Despite the eradication of polio and measles being within reach,²⁵ the research output on these vaccines remained relatively modest throughout the study period. Moreover, notwithstanding the rise in public vaccine hesitancy in recent years,^{26 27} it did not intensify as much as expected after the COVID-19 pandemic, proving vulnerable to sudden disruptions. In contrast, we observed higher-than-expected post-pandemic publication volumes regarding the influenza vaccine, likely due to its analogies with the SARS-CoV-2 vaccine. These findings align with abundant evidence on the negative impact of the SARS-CoV-2 pandemic on routine childhood vaccinations, both in terms of vaccine uptake^{28 29} and public trust³⁰, and may be attributed to the temporary redirection of academic interest and research funding towards COVID-19-related research. The influence of SARS-CoV-2 on PSV research was further highlighted by a post-pandemic increase in articles on other primary vaccines mentioning the SARS-CoV-2 vaccine.

This pattern, especially striking for vaccines that share few or no similarities with the SARS-CoV-2 vaccine, likely reflects a momentary shift of the global health research agenda due to the very peculiar and worrisome situation of the pandemic threat. However, it also suggests that PSV research is highly sensitive to time-dependent and context-dependent funding and publication opportunities, which in turn influence researchers’ actions.¹⁶ This result underscores the importance of maintaining sustained academic attention on long-standing research objectives, such as vaccine-preventable diseases elimination and eradication efforts.

Our statistical analyses of geographical, economic and epidemiological factors consistently showed that country income levels critically shape the landscape of PSV research, inducing an economically driven imbalance in both the global distribution and direction of research efforts.

In agreement with previous literature,^{8 10–12} we found that HIC produced the majority of global PSV research, typically redirecting their research on polio and measles towards LMIC and, on the other hand, attracting research efforts on HPV and influenza vaccines.

Our regression analysis also suggested a gradual shift in academic attention from LMIC to HIC, particularly for measles, but also for HPV and influenza. We showed that such a trend may be partially explained by changes in the epidemiology of the related diseases, although other factors likely played a significant role as well. These may include a country’s historical experience with the disease, the presence of local outbreaks, local vaccination recommendations and policies, political context, an unequal

distribution of research funding, global publication bias and countries' involvement in international public health endeavours, such as disease elimination initiatives. These results also suggest that the gap left by shifting attention to HIC is not currently filled with domestic research in LMIC.

Income-dependent disparities were also evident through the disproportionate amount of foreign authorship in studies concerning LMIC, especially in PSV research on polio and measles. This suggests a structural involvement of external researchers in LMIC, even in cases concerning very local health challenges. Except for polio research, where the historical imbalances persist, foreign research contributions became more evenly distributed across income groups for all vaccines, indicating an encouraging increase in academic cross-income-group collaborations. Combining this result with an overall increased attention to higher-income countries, however, our analyses also suggest that the levelling of foreign efforts might be driven by increased collaborations among HIC, likely at the expense of lower-income ones, where such research is often most needed.

Overall, our regression analyses provided strong evidence that a country's lower income significantly limits its capacity to perform domestic research, even in areas with high disease incidence, resulting in an inequitable and likely inefficient reliance on external research activities to address critical local health challenges.

We acknowledge some limitations. First, our literature search strategy primarily employed English-language keywords and a substantial portion of non-English scientific literature may not be indexed in the two databases used. As a result, we may have missed relevant studies published in other languages lacking an English abstract. Because English-language publications are not fully representative of the global scientific landscape, this may have introduced a language bias into our literature corpus. However, the vast majority of global scientific output is currently published in English.³¹ Hence, any language bias is likely to be limited in scope and not expected to substantially affect the main patterns identified in the analysis.

Second, although our ML-assisted article selection achieved strong performance metrics, some misclassification may have occurred, potentially leading to the inclusion of a few irrelevant articles or the exclusion of some pertinent ones. Yet, given the scale of the dataset and the inherent subjectivity in relevance judgements, even a fully manual screening process would likely have introduced comparable inconsistencies. Importantly, the volume and the consistency of the retained articles still provide a reliable basis for mapping broader scientific trends.

Lastly, the inclusion of additional contextual factors that may shape global attention on PSV research could offer deeper explanatory power. However, in order to ensure complete, consistent and reliable cross-country comparison on a global scale, we chose to leverage only

indicators provided by internationally recognised organisations. Collating national datasets may have introduced uncertainty and unknown sources of potential bias, ultimately compromising the robustness and comparability of our analysis.

CONCLUSIONS

The persistent misalignment between countries with the financial capacity to conduct PSV research and those most in need of its outcomes highlights entrenched inequities within this field. These disparities are not unique to PSV research but are rather symptomatic of broader structural imbalances that pervade global health research. Research priorities—particularly in HIC—are often shaped by internal agendas and funding opportunities rather than global need. This hampers the potential impact of research efforts in achieving broader health outcomes, such as increased vaccine uptake in underserved settings.³²

Our findings suggest that such imbalances are likely to be exacerbated during periods of global disruptions, as demonstrated by the COVID-19 pandemic, with potentially detrimental consequences for both ongoing public health interventions and future research directions. By providing a clear view of the main drivers of PSV research and revealing the risks of shifting academic focus, this study highlights the need to decolonise PSV research and, broadly, global health research. Addressing historical imbalances requires promoting equitable, inclusive research partnerships, and redistributing leadership, ownership and priority-setting power across the global research community. Key principles for establishing equitable research dynamics include sustainability, policy-oriented objectives, transparency, equitable data access and meaningful community engagement.^{33–37} Existing ethical guidance can help researchers design projects in alignment with these values,³⁸ and funding mechanisms should embed strong incentives to promote such practices.³⁹ Future funding allocation criteria should prioritise research with the highest potential health impact for vulnerable populations—such as improving vaccine coverage among underimmunised children—over purely academic or innovation-driven considerations.⁴⁰ Grants must be accessible to institutions in LMIC directly, fostering application diversity and enhancing local research capacity.^{3,41} Achieving key global health targets, such as the eradication of polio and measles, requires sustainable investments guided by clear long-term strategies and implemented within well-defined timeframes, particularly in low-resource settings.⁴² Currently, decision-making power remains highly concentrated in a small group of actors, mostly from HIC, yet continued support may be subject to political or economic fluctuations. This can lead to funding disruptions with potentially serious implications for the populations affected. A sustainable transition towards increased domestic investment in health research is therefore critical and must include

robust exit strategies, early planning, stakeholder engagement and dedicated pre-transition resources.^{3 43 44} Scientific journals are also playing a crucial role in reshaping inequitable research practices by establishing systems to identify and discourage submissions that violate equity principles related to content and authorship.^{45 46} Our work highlights the need for a systemic change across funding structures, authorship norms and publication practices in order to redress long-standing inequities in global health research and to ensure that research efforts ultimately benefit the most affected.⁴⁷

Author affiliations

¹Dondena Centre for Research on Social Dynamics and Public Policy, Bocconi University, Milano, Italy

²BIDSA Bocconi Institute for Data Science and Analytics, Bocconi University, Milano, Italy

³Nuffield College, University of Oxford, Oxford, UK

⁴Department of Economics and Management, University of Pisa, Pisa, Italy

⁵Department of Statistics, Informatics, and Applications "Giuseppe Parenti", University of Florence, Florence, Italy

⁶Department of Computer Science, Christian-Albrechts-Universität zu Kiel, Kiel, Germany

⁷CERGAS Centre for Research on Health and Social Care Management, Bocconi University, Milano, Italy

⁸CMCC Foundation, Euro-Mediterranean Center on Climate Change, Milan, Italy

⁹RFF-CMCC European Institute on Economics and the Environment, Milan, Italy

¹⁰Centre for Computational Intelligence, School of Computer Science and Electronic Engineering, University of Essex, Colchester, UK

¹¹Smart Health Technologies Group, Institute of Public Health and Wellbeing, University of Essex, Colchester, UK

Contributors DB, VO, ZA, CC, LPL, AT, SS, JA-P and AM contributed to the conceptualisation by formulating the research idea, objectives and research questions. VO, LPL, ED'A and AM developed the search string, screening strategy and annotated the dataset for relevance. DS and JA-P collected and curated the dataset, while DB, VO, ZA and CC enriched it with relevant attributes and performed data cleaning. DB, VO, ZA, CC and AM designed the study methods and performed technical analyses, including machine-assisted literature searches, enrichment via natural language processing, network analysis and other formal analyses. CC and ZA conducted the regression analysis. DB, VO, ZA, CC and AM produced the results and drafted the initial manuscript. All authors interpreted the results, reviewed and edited the drafts. DB and VO led and coordinated the project, AM supervised and secured funding. AM is the guarantor. Beyond the AI techniques described in the Methods section of the manuscript, we used AI to shorten and improve the writing style of individual sentences.

Funding Supported by the ERC Grant 'IMMUNE' (ID: 101003183), PNRR Grant BEHAVE-MOD (ID: SA-02. P0001) and the Bocconi Covid Crisis Lab, supported by the 'Fondazione Romeo ed Enrica Invernizzi'.

Map disclaimer The depiction of boundaries on this map does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. This map is provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study did not include human participants and is thus exempt from institutional review board approval and the requirement for informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The database containing all scientific publications included in this analysis will be made available on reasonable request for scientific research purposes. For further information or to request access, please contact the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Vittoria Offeddu <https://orcid.org/0000-0003-4381-7689>

Soheil Shayegh <https://orcid.org/0000-0002-8960-8244>

REFERENCES

- Koplan JP, Bond TC, Merson MH, *et al*. Towards a common definition of global health. *The Lancet* 2009;373:1993–5.
- Smeeth L, Kyobutungi C. Reclaiming global health. *The Lancet* 2023;401:625–7.
- Charani E, Abimbola S, Pai M, *et al*. Funders: The missing link in equitable global health research? *PLOS Glob Public Health* 2022;2:e0000583.
- Cash-Gibson L, Rojas-Gualdrón DF, Pericàs JM, *et al*. Inequalities in global health inequalities research: A 50-year bibliometric analysis (1966–2015). *PLoS ONE* 2018;13:e0191901.
- Global Forum for Health Research & World Health Organization. The 10/90 (ten ninety) report on health research 2003–2004, 2004. Available: <https://iris.who.int/handle/10665/44386> [Accessed 16 Jul 2024].
- The WHO Global Health Observatory. IA2030 IG 2.1: Number of zero dose children, 2024. Available: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/7792> [Accessed 19 Sep 2024].
- de Figueiredo A, Simas C, Karafillakis E, *et al*. Mapping global trends in vaccine confidence and investigating barriers to vaccine uptake: a large-scale retrospective temporal modelling study. *The Lancet* 2020;396:898–908.
- Sweileh WM. Bibliometric analysis of global scientific literature on vaccine hesitancy in peer-reviewed journals (1990–2019). *BMC Public Health* 2020;20:1–15.
- Fernandes S, Jit M, Bozzani F, *et al*. A bibliometric analysis of systematic reviews on vaccines and immunisation. *Vaccine (Auckl)* 2018;36:2254–61.
- Chen C, Yang Q, Tian H, *et al*. Bibliometric and visual analysis of vaccination hesitancy research from 2013 to 2022. *Hum Vaccin Immunother* 2023;19:2226584.
- Sweileh WM. Bibliometric analysis of scientific research publications on COVID-19 vaccine hesitancy (2020 – 2021). *Electronic Journal of General Medicine* 2023;20:em500.
- Jaca A, Iwu-Jaja CJ, Balakrishna Y, *et al*. A global bibliometric analysis of research productivity on vaccine hesitancy from 1974 to 2019. *Hum Vaccin Immunother* 2021;17:3016–22.
- Emery N, Dugeril A, Flahault A. Vaccine Hesitancies Across the World in the Era of COVID-19. *Am J Public Health* 2022;112:1579–81.
- He J, Liu X, Lu X, *et al*. The impact of COVID-19 on global health journals: an analysis of impact factor and publication trends. *BMJ Glob Health* 2023;8:e011514.
- Riccaboni M, Verginer L. The impact of the COVID-19 pandemic on scientific research in the life sciences. *PLoS ONE* 2022;17:e0263001.
- Aviv-Reuven S, Rosenfeld A. Publication patterns' changes due to the COVID-19 pandemic: a longitudinal and short-term scientometric analysis. *Scientometrics* 2021;126:6761–84.
- Bramer WM, De Jonge GB, Rethlefsen ML, *et al*. A systematic approach to searching: an efficient and complete method to develop literature searches. *jmla* 2018;106.
- Bramer WM, Rethlefsen ML, Kleijnen J, *et al*. Optimal database combinations for literature searches in systematic reviews: a prospective exploratory study. *Syst Rev* 2017;6:245.

- 19 Chicco D, Jurman G. The advantages of the Matthews correlation coefficient (MCC) over F1 score and accuracy in binary classification evaluation. *BMC Genomics* 2020;21:6.
- 20 World Bank Group. World Bank Country and Lending Groups. 2024.
- 21 World Health Organization. Polio (Pol3) Immunization coverage estimates by country, Available: <https://apps.who.int/gho/data/node.main.A831> [Accessed 15 Jul 2024].
- 22 Measles-containing-vaccine second-dose (mcv2) immunization coverage by the locally recommended age (%). 2024. Available: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/measles-containing-vaccine-second-dose-\(mcv2\)-immunization-coverage-by-the-nationally-recommended-age-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/measles-containing-vaccine-second-dose-(mcv2)-immunization-coverage-by-the-nationally-recommended-age-(-)) [Accessed 15 Jul 2024].
- 23 World Health Organization. Poliomyelitis - number of reported cases, 2024. Available: <https://www.who.int/data/gho/data/indicators/indicator-details/GHO/poliomyelitis---number-of-reported-cases> [Accessed 15 Jul 2024].
- 24 World Health Organization. Measles - number of reported cases, 2024. Available: <https://www.who.int/data/gho/data/indicators/indicator-details/GHO/measles---number-of-reported-cases> [Accessed 15 Jul 2024].
- 25 Higgins DM, O'Leary ST. A World without Measles and Rubella: Addressing the Challenge of Vaccine Hesitancy. *Vaccines (Basel)* 2024;12:694.
- 26 Pandey A, Galvani AP. Exacerbation of measles mortality by vaccine hesitancy worldwide. *Lancet Glob Health* 2023;11:e478–9.
- 27 Harvard International Review. Polio in a post vaccine world the consequences of anti-vaccination sentiments. 2023. Available: <https://hir.harvard.edu/polio-in-a-post-vaccine-world-the-consequences-of-anti-vaccination-sentiments/>
- 28 Shet A, Carr K, Danovaro-Holliday MC, et al. Impact of the SARS-CoV-2 pandemic on routine immunisation services: evidence of disruption and recovery from 170 countries and territories. *Lancet Glob Health* 2022;10:e186–94.
- 29 Lassi ZS, Naseem R, Salam RA, et al. The Impact of the COVID-19 Pandemic on Immunization Campaigns and Programs: A Systematic Review. *Int J Environ Res Public Health* 2021;18:988:1–19.
- 30 Lazarus JV, White TM, Wyka K, et al. Influence of COVID-19 on trust in routine immunization, health information sources and pandemic preparedness in 23 countries in 2023. *Nat Med* 2024;30:1559–63.
- 31 Scientific publishing has a language problem. *Nat Hum Behav* 2023;7:1019–20.
- 32 Agnandji ST, Tsassa V, Conzelmann C, et al. Patterns of biomedical science production in a sub-Saharan research center. *BMC Med Ethics* 2012;13.
- 33 Monette EM, McHugh D, Smith MJ, et al. Informing “good” global health research partnerships: A scoping review of guiding principles. *Glob Health Action* 2021;14:1892308.
- 34 Evertsz N, Bull S, Pratt B. What constitutes equitable data sharing in global health research? A scoping review of the literature on low-income and middle-income country stakeholders' perspectives. *BMJ Glob Health* 2023;8:e010157:1–11.
- 35 Achana F, Boah M, Chirawurah D, et al. Equitable global health research collaborations with a mind of human dignity. *Front Educ* 2023;8:1–5.
- 36 Hodson DZ, Etoundi YM, Parikh S, et al. Striving towards true equity in global health: A checklist for bilateral research partnerships. *PLOS Glob Public Health* 2023;3:e0001418.
- 37 Saleh S, Masekela R, Heinz E, et al. Equity in global health research: A proposal to adopt author reflexivity statements. *PLOS Glob Public Health* 2022;2:e0000160.
- 38 Pratt B. Research for health justice framework. 2025. Available: <https://researchforhealthjustice.squarespace.com/general-2> [Accessed 7 May 2025].
- 39 Pratt B, Hyder AA. Designing research funding schemes to promote global health equity: An exploration of current practice in health systems research. *Dev World Bioeth* 2018;18:76–90.
- 40 Murray M, Mubiligi J. An Approach to Building Equitable Global Health Research Collaborations. *Ann Glob Health* 2020;86:126.
- 41 Chikwari CD, Tadesse AW, Shanaube K, et al. Achieving equitable leadership in Global Health partnerships: barriers experienced and strategies to improve grant funding for early- and mid-career researchers. *BMC Glob Public Health* 2024;2:17.
- 42 Stenberg K, Hanssen O, Edejer T-T, et al. Financing transformative health systems towards achievement of the health Sustainable Development Goals: a model for projected resource needs in 67 low-income and middle-income countries. *Lancet Glob Health* 2017;5:e875–87.
- 43 Olufadewa I, Adesina M, Ayorinde T. Global health in low-income and middle-income countries: a framework for action. *Lancet Glob Health* 2021;9:e899–900.
- 44 Huffstetler HE, Bandara S, Bharali I, et al. The impacts of donor transitions on health systems in middle-income countries: a scoping review. *Health Policy Plan* 2022;37:1188–202.
- 45 Khan SA. Decolonising global health by decolonising academic publishing. *BMJ Glob Health* 2022;7:1–6.
- 46 Olusanya BO, Mallewa M, Ogbo FA. Beyond pledges: academic journals in high-income countries can do more to decolonise global health. *BMJ Glob Health* 2021;6:1–3.
- 47 Büyüm AM, Kenney C, Koris A, et al. Decolonising global health: if not now, when? *BMJ Glob Health* 2020;5:e003394:1–4.